CALCULUS, Fourth Edition, Hughes-Hallett, Gleason, McCallum, et al.

## Chapter 2

Section Suggested Problems

- 2.1 1, 3, 5-8, 11, 12, 14, 15.
- 2.2 1, 3, 5, 8-13, 15, 16, 21.
- 2.3 1-13 odd, 17, 19, 33, 36, 40-42.
- 2.4 1, 3, 6, 7, 10, 11, 14, 15, 17.
- 2.5 1-5, 7, 9, 11, 14, 15, 18 23.
- 2.6 1-5, 6, 10, 15.

Review 7, 8, 19, 20, 28.




Figure 1:  $y = ax^2 + bx + c$ 

Please recall Problem 46 on page 60 and what we did in class related to this problem. Here we replace the function  $y = x^2$  with a general parabola  $y = ax^2 + bx + c$  and repeat what we did in class to find the slope of the tangent line to this parabola at the point  $(x_0, y_0)$  where  $y_0 = ax_0^2 + bx_0 + c$ . Let  $(0, \beta)$  be a point on y-axes. The slope of the line through the points  $(0, \beta)$  and  $(x_0, y_0)$  is

$$\frac{y_0 - \beta}{x_0 - 0} = \frac{ax_0^2 + bx_0 + c - \beta}{x_0}.$$

The equation of the line through the points  $(0,\beta)$  and  $(x_0,y_0)$  is

$$y = \frac{ax_0^2 + bx_0 + c - \beta}{x_0}x + \beta.$$

To find where this line intersects the parabola  $y = ax^2 + bx + c$  we solve for x the equation:

$$ax^{2} + bx + c = \frac{ax_{0}^{2} + bx_{0} + c - \beta}{x_{0}}x + \beta.$$

Multiplying through by  $x_0$  we get

$$ax_0x^2 + bx_0x + cx_0 = (ax_0^2 + bx_0 + c - \beta)x + \beta x_0.$$

Grouping the terms in the last equation and factoring gives

$$0 = ax_0x^2 + bx_0x + cx_0 - axx_0^2 - bxx_0 - cx + \beta x - \beta x_0$$
  
=  $ax_0x(x - x_0) - c(x - x_0) + \beta(x - x_0)$   
=  $(ax_0x - c + \beta)(x - x_0)$ .

Hence the x-coordinates of the points of intersection of the line and the parabola are

$$x = x_1 = \frac{c - \beta}{ax_0}$$
 and  $x = x_0$ .

These two points coincide if  $c - \beta = ax_0^2$ , that is if  $\beta = c - ax_0^2$ . In this case the line is the tangent line to the parabola at the point  $(x_0, y_0)$  and the slope of the tangent line is

$$\frac{ax_0^2 + bx_0 + c - \beta}{x_0} = \frac{ax_0^2 + bx_0 + c - c + ax_0^2}{x_0} = \frac{2ax_0^2 + bx_0}{x_0} = 2ax_0 + b.$$

Thus the slope of the tangent line to the parabola  $y = ax^2 + bx + c$  at the point  $(x_0, y_0)$  where  $y_0 = ax_0^2 + bx_0 + c$  is given by the formula  $2ax_0 + b$ .

In Figure 1 I used  $a = -2, b = 5, c = -1, x_0 = 2$  and  $\beta = 3$ . What is the slope of the tangent line to this parabola at the point (2,1)? (Notice that the horizontal and vertical units are different.)