

# Jordan normal form

Branko Ćurgus

December 5, 2013 at 13:04

Throughout this note  $\mathcal{V}$  is a nontrivial finite dimensional vector space over  $\mathbb{C}$ . We set  $n = \dim \mathcal{V}$ . The symbol  $\mathbb{N}$  denotes the set of positive integers and  $i, j, k, l, m, n, p, q, r \in \mathbb{N}$ . For  $T \in \mathcal{L}(\mathcal{V})$  by  $\mathcal{N}(T)$  we denote the null-space and by  $\mathcal{R}(T)$  the range of  $T$ .

## 1 Nilpotent operators

An operator  $N \in \mathcal{L}(\mathcal{V})$  is *nilpotent* if there exists  $q \in \mathbb{N}$  such that  $N^q = 0$ . If  $N^q = 0$  and  $N^{q-1} \neq 0$ , then  $q$  is called *the degree of nilpotency* of  $N$ .

It is not difficult to show that in a finite dimensional space  $\mathcal{V}$  the degree of nilpotency  $q$  satisfies  $q \leq \dim \mathcal{V}$ .

tjbn

**Theorem 1.1.** *Let  $\mathcal{V}$  be a nontrivial finite dimensional vector space over  $\mathbb{C}$  with  $n = \dim \mathcal{V}$ . Let  $N \in \mathcal{L}(\mathcal{V})$  be a nilpotent operator such that  $m = \dim \mathcal{N}(N)$ . Then there exist vectors  $v_1, \dots, v_m \in \mathcal{V}$  and positive integers  $q_1, \dots, q_m$  such that the vectors*

$$N^{q_k-1}v_k, \quad k \in \{1, \dots, m\},$$

*form a basis of  $\mathcal{N}(N)$  and the vectors*

$$v_k, Nv_k, \dots, N^{q_k-1}v_k, \quad k \in \{1, \dots, m\},$$

*form a basis of  $\mathcal{V}$ .*

*Proof.* The proof is by induction on the dimension  $n$ . For  $n = 1$ ,  $N^{dim}(V) = N^1 = 0$ , so the statement is trivially true for  $n = 1$ . Let  $n \in \mathbb{N}$  and assume that the statement is true for any vector space of dimension less or equal to  $n$ . It is always a good idea to be specific and state what is being assumed. Let  $n \in \mathbb{N}$  be such that  $n > 1$ . The following implication is our inductive hypothesis:

If  $\mathcal{W}$  is a vector space over  $\mathbb{C}$  such that  $\dim \mathcal{W} < n$  and if  $M \in \mathcal{L}(\mathcal{W})$  is a nilpotent operator such that  $l = \dim \mathcal{N}(M)$ , then there exist  $w_1, \dots, w_l \in \mathcal{W}$  and positive integers  $p_1, \dots, p_l$  such that the vectors

$$M^{p_j-1}w_j, \quad j \in \{1, \dots, l\},$$

form a basis of  $\mathcal{N}(M)$  and the vectors

$$w_j, Mw_j, \dots, M^{p_j-1}w_j, \quad j \in \{1, \dots, l\},$$

form a basis of  $\mathcal{W}$ .

Next we present a proof of the inductive step.

Let  $\mathcal{V}$  be a nontrivial finite dimensional vector space over  $\mathbb{C}$  with  $\dim \mathcal{V} = n$ . Let  $N \in \mathcal{L}(\mathcal{V})$  be a nilpotent operator.

First notice that if  $N = 0$ , then  $\mathcal{N}(N) = \mathcal{V}$  and the claim is trivially true. In this case  $m = n$  and any basis  $v_1, \dots, v_n$  of  $\mathcal{V}$  with positive integers  $q_1 = \dots = q_n = 1$  satisfies the requirement of the theorem. From now on we assume that  $N \neq 0$ .

Set  $m = \dim \mathcal{N}(N)$  and  $\mathcal{W} = \mathcal{R}(N)$ . Since all powers of an invertible operator are invertible and a power of  $N$  is 0,  $N$  is not invertible. Thus  $m = \dim \mathcal{N}(N) \geq 1$ . By the famous ‘‘rank-nullity’’ theorem  $\dim \mathcal{W} < n$ . Since  $N \neq 0$ ,  $\dim \mathcal{W} > 0$ . It is clear that  $\mathcal{W}$  is invariant under  $N$ . Set  $M \in \mathcal{L}(\mathcal{W})$  to be the restriction of  $N$  onto  $\mathcal{W}$ . Then  $M \in \mathcal{L}(\mathcal{W})$ . Since  $N$  is nilpotent,  $M$  is nilpotent as well. Clearly,  $\mathcal{N}(M) = \mathcal{N}(N) \cap \mathcal{R}(N)$ . Set  $l = \dim \mathcal{N}(M)$ . The vector space  $\mathcal{W}$  and the operator  $M$  satisfy all the assumptions of the inductive hypothesis. This allows us to deduce that there exist  $w_1, \dots, w_l \in \mathcal{W}$  and positive integers  $p_1, \dots, p_l$  such that the vectors

$$M^{p_j-1}w_j, \quad j \in \{1, \dots, l\}, \quad (1) \quad \boxed{\text{eqbaNM}}$$

form a basis of  $\mathcal{N}(M) = \mathcal{N}(N) \cap \mathcal{R}(N)$  and the vectors

$$w_j, Mw_j, \dots, M^{p_j-1}w_j, \quad j \in \{1, \dots, l\}, \quad (2) \quad \boxed{\text{eqbaW}}$$

form a basis of  $\mathcal{W} = \mathcal{R}(N)$ . Since  $w_j \in \mathcal{R}(N)$ , there exist  $v_j \in \mathcal{V}$  such that  $w_j = Nv_j$  for all  $j \in \{1, \dots, l\}$ . We know from (1) that the vectors

$$M^{p_1-1}w_1 = N^{p_1}v_1, \dots, M^{p_l-1}w_l = N^{p_l}v_l,$$

form a basis of  $\mathcal{N}(M) = \mathcal{N}(N) \cap \mathcal{R}(N)$ . Recall that  $m = \dim \mathcal{N}(N)$  and  $l \leq m$ . Let  $v_{l+1}, \dots, v_m$  be such that

$$N^{p_1}v_1, \dots, N^{p_l}v_l, v_{l+1}, \dots, v_m, \quad (3) \quad \boxed{\text{eqbaNN}}$$

form a basis of  $\mathcal{N}(N)$ . (It is possible that  $l = m$ . In this case we already have a basis of  $\mathcal{N}(N)$  and the last step can be skipped.)

Now let us review the stage: We started with the basis

$$w_j = Nv_j, Mw_j = N^2v_j, \dots, M^{p_j-1}w_j = N^{p_j}v_j, \quad j \in \{1, \dots, l\},$$

of  $\mathcal{W} = \mathcal{R}(N)$ , with  $\dim \mathcal{R}(N)$  vectors to which we add the vectors  $v_1, \dots, v_m$ . Now we have  $m + \dim \mathcal{R}(N) = \dim \mathcal{N}(N) + \dim \mathcal{R}(N) = \dim \mathcal{V} = n$  vectors:

$$v_j, Nv_j, N^2v_j, \dots, N^{p_j}v_j, \quad j \in \{1, \dots, l\}, \quad v_{l+1}, \dots, v_m. \quad (4) \quad \boxed{\text{eqbaV}}$$

For easier writing set

$$q_k = \begin{cases} p_k + 1 & \text{if } k \in \{1, \dots, l\} \\ 1 & \text{if } k \in \{l+1, \dots, m\}. \end{cases}$$

Then (4) can be rewritten as

$$v_k, Nv_k, N^2v_k, \dots, N^{q_k-1}v_k, \quad k \in \{1, \dots, m\}. \quad (5) \quad \boxed{\text{eqbaV1}}$$

Next we will prove that the vectors in (5) are linearly independent. Let  $\alpha_{k,j} \in \mathbb{C}, j \in \{0, \dots, q_k - 1\}, k \in \{1, \dots, m\}$  be such that

$$\sum_{k=1}^m \sum_{j=0}^{q_k-1} \alpha_{k,j} N^j v_k = 0. \quad (6) \quad \boxed{\text{eqlin}}$$

Applying  $N$  to the last equality yields

$$\sum_{k=1}^l \sum_{j=0}^{q_k-2} \alpha_{k,j} N^{j+1} v_k = \sum_{k=1}^l \sum_{j=0}^{p_k-1} \alpha_{k,j} M^j w_k = 0.$$

Since the vectors in the last double sum are the vectors from (2) they are linearly independent. Therefore

$$\alpha_{k,0} = \dots = \alpha_{k,q_k-2} = 0, \quad k \in \{1, \dots, l\}.$$

Substituting these values in (6) we get

$$\sum_{k=1}^m \alpha_{k,q_k-1} N^{q_k-1} v_k = 0.$$

But, beautifully, the vectors in the last sum are exactly the vectors in (3) which are linearly independent. Thus

$$\alpha_{k,q_k-1} = 0, \quad k \in \{1, \dots, m\}.$$

This completes the proof that all the coefficients in (6) must be zero. Thus, the vectors in (5) are linearly independent. Since there are exactly  $n$  vectors in (5) they form a basis of  $\mathcal{V}$ . This completes the proof.  $\square$

## 2 A Decomposition of a Vector Space

$\boxed{\text{li}}$  **Lemma 2.1.** *Let  $\mathcal{V}$  be a vector space over  $\mathbb{F}$ . Let  $A$  and  $B$  be commuting linear operators on  $\mathcal{V}$ . Then  $\mathcal{N}(B)$  and  $\mathcal{R}(B)$  are invariant subspaces for  $A$ .*

*Proof.* This is a very simple exercise.  $\square$

$\boxed{\text{pges}}$  **Proposition 2.2.** *Let  $\mathcal{V}$  be a vector space over  $\mathbb{F}$ . Let  $T \in \mathcal{L}(\mathcal{V})$ . If  $\lambda$  and  $\mu$  are distinct eigenvalues of  $T$  and  $j$  and  $k$  are natural numbers, then*

$$\mathcal{N}((T - \lambda I)^j) \cap \mathcal{N}((T - \mu I)^k) = \{0_{\mathcal{V}}\}.$$

*Proof.* The set equality in the proposition is equivalent to the implication

$$v \in \mathcal{N}((T - \mu I)^k) \setminus \{0_{\mathcal{V}}\} \Rightarrow v \notin \mathcal{N}((T - \lambda I)^j).$$

We will prove this implication. Let  $v \in \mathcal{V}$  be such that  $(T - \mu I)^k v = 0_{\mathcal{V}}$  and  $v \neq 0_{\mathcal{V}}$ . Let  $i \in \{1, \dots, k\}$  be such that  $(T - \mu I)^i v = 0_{\mathcal{V}}$  and  $(T - \mu I)^{i-1} v \neq 0_{\mathcal{V}}$ . Set  $w = (T - \mu I)^{i-1} v$ . Then  $w$  is an eigenvector of  $T$  corresponding to  $\mu$ , that is  $Tw = \mu w$  and  $w \neq 0$ . Then (as we must have proven before), for an arbitrary polynomial  $p \in \mathbb{C}[z]$  we have  $p(T)w = p(\mu)w$ . In particular

$$(T - \lambda I)^l w = (\mu - \lambda)^l w \quad \text{for all } l \in \mathbb{N}.$$

Since  $\mu - \lambda \neq 0$  and  $w \neq 0_{\mathcal{V}}$  we have that

$$(T - \lambda I)^l w \neq 0_{\mathcal{V}} \quad \text{for all } l \in \mathbb{N}.$$

Consequently,

$$(T - \lambda I)^l (T - \mu I)^{i-1} v \neq 0_{\mathcal{V}} \quad \text{for all } l \in \mathbb{N}.$$

Since the operators  $(T - \lambda I)^l$  and  $(T - \mu I)^{i-1}$  commute we have

$$(T - \mu I)^{i-1} (T - \lambda I)^l v \neq 0_{\mathcal{V}} \quad \text{for all } l \in \mathbb{N}.$$

Therefore  $(T - \lambda I)^l v \neq 0_{\mathcal{V}}$  for all  $l \in \mathbb{N}$ . Hence  $v \notin \mathcal{N}((T - \lambda I)^j)$ . This proves the proposition.  $\square$

cges

**Corollary 2.3.** *Let  $\mathcal{V}$  be a finite dimensional vector space over  $\mathbb{F}$ . Let  $T \in \mathcal{L}(\mathcal{V})$ . If  $\lambda$  and  $\mu$  are distinct eigenvalues of  $T$  and  $j$  and  $k$  are natural numbers, then*

$$\mathcal{N}((T - \lambda I)^j) \subseteq \mathcal{R}((T - \mu I)^k).$$

*Proof.* Since the operators  $(T - \lambda I)^j$  and  $(T - \mu I)^k$  commute, by Lemma 2.1,  $\mathcal{N}((T - \lambda I)^j)$  is invariant under  $(T - \mu I)^k$ . Denote by  $S$  the restriction of  $(T - \mu I)^k$  onto  $\mathcal{N}((T - \lambda I)^j)$ . Since clearly,

$$\mathcal{N}(S) = \mathcal{N}((T - \lambda I)^j) \cap \mathcal{N}((T - \mu I)^k).$$

Proposition 2.2 implies that  $S$  is an injection, and thus bijection. Hence,

$$S\left(\mathcal{N}((T - \lambda I)^j)\right) = \mathcal{N}((T - \lambda I)^j)$$

and consequently

$$\mathcal{N}((T - \lambda I)^j) = (T - \mu I)^k\left(\mathcal{N}((T - \lambda I)^j)\right) \subseteq \mathcal{R}((T - \mu I)^k).$$

$\square$

le-red

**Lemma 2.4.** *Let  $\mathcal{V}$  be a vector space over  $\mathbb{F}$ . Let  $\mathcal{U}$  and  $\mathcal{W}$  be subspaces of  $\mathcal{V}$  such that*

$$\mathcal{V} = \mathcal{U} \oplus \mathcal{W}.$$

*Let  $S \in \mathcal{L}(V)$  be such that  $S\mathcal{U} \subseteq \mathcal{U}$  and  $S\mathcal{W} \subseteq \mathcal{W}$ . If  $\mathcal{N}(S) \cap \mathcal{W} = \{0\}$ , then*

$$\mathcal{N}((S|_{\mathcal{U}})^j) = \mathcal{N}(S^j) \quad \text{for all } j \in \mathbb{N}. \quad (7)$$

eq-ns=ns

*Proof.* Assume  $\mathcal{N}(S) \cap \mathcal{W} = \{0\}$ . We first prove the equality for  $j = 1$ . Since  $\mathcal{N}(S|_{\mathcal{U}}) = \mathcal{N}(S) \cap \mathcal{U}$ , the inclusion  $\mathcal{N}(S|_{\mathcal{U}}) \subseteq \mathcal{N}(S)$  is clear. Let  $v \in \mathcal{N}(S)$  be arbitrary. Then  $v = u + w$  with  $u \in \mathcal{U}$  and  $w \in \mathcal{W}$ . Applying  $S$  to this identity we get  $0 = Sv = Su + Sw$ . Since  $Su \in \mathcal{U}$  and  $Sw \in \mathcal{W}$ , the assumption that the sum of  $\mathcal{U}$  and  $\mathcal{W}$  is direct yields  $Sw = 0$ . Since  $\mathcal{N}(S) \cap \mathcal{W} = \{0\}$ , we have  $w = 0$ . Thus,  $v \in \mathcal{U}$ , and hence  $v \in \mathcal{N}(S|_{\mathcal{U}})$ .

To prove (7) for arbitrary  $j \in \mathbb{N}$  we will first prove that

$$\mathcal{N}(S^j) \cap \mathcal{W} = \{0\} \quad \text{for all } j \in \mathbb{N}. \quad (8)$$

eq-nul-j

A simple proof proceeds by mathematical induction. The statement in (8) is true for  $j = 1$ . Let  $j \in \mathbb{N}$  and assume that the statement in (8) is true for  $j$ . Now assume that  $w \in \mathcal{W}$  and  $S^{j+1}w = 0$ . Then  $Sw \in \mathcal{W}$  and  $S^j(Sw) = 0$ . By the inductive hypothesis, that is  $\mathcal{N}(S^j) \cap \mathcal{W} = \{0\}$  we conclude  $Sw = 0$ . Since  $\mathcal{N}(S) \cap \mathcal{W} = \{0\}$ , we deduce that  $w = 0$ .

Having (8), we can apply the equality proved in the first part of the proof to the operator  $S^j$ .  $\square$

co-redsp

**Corollary 2.5.** *Let  $\mathcal{V}$  be a finite dimensional vector space over  $\mathbb{F}$ . Let  $\mathcal{U}$  and  $\mathcal{W}$  be subspaces of  $\mathcal{V}$  such that*

$$\mathcal{V} = \mathcal{U} \oplus \mathcal{W}.$$

Let  $T \in \mathcal{L}(V)$  be such that  $T\mathcal{U} \subseteq \mathcal{U}$  and  $T\mathcal{W} \subseteq \mathcal{W}$ . Then

$$\sigma(T|_{\mathcal{U}}) \cup \sigma(T|_{\mathcal{W}}) = \sigma(T). \quad (9)$$

eq-sp-par

If  $\lambda \in \sigma(T)$  and  $\lambda \notin \sigma(T|_{\mathcal{W}})$ , then  $\lambda \in \sigma(T|_{\mathcal{U}})$  and

$$\mathcal{N}((T|_{\mathcal{U}} - \lambda I)^j) = \mathcal{N}((T - \lambda I)^j) \quad \text{for all } j \in \mathbb{N}. \quad (10)$$

eq-gen-es=

*Proof.* The inclusion  $\subseteq$  in (9) is clear. To prove  $\supseteq$ , let  $\lambda \in \sigma(T)$  and let  $v \neq 0$  be such that  $Tv = \lambda v$ . Let  $v = u + w$ , with  $u \in \mathcal{U}$  and  $w \in \mathcal{W}$ . Since  $v \neq 0$  we have  $u \neq 0$  or  $w \neq 0$ . Applying  $T$  to both sides of  $v = u + w$  and using the fact that  $v$  is an eigenvalue corresponding to  $\lambda$  we get  $Tu + Tw = Tv = \lambda v = \lambda u + \lambda w$ . Consequently,  $(Tu - \lambda u) + (Tw - \lambda w) = 0$ . Since the sum  $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$  is direct and  $Tu - \lambda u \in \mathcal{U}$  and  $Tw - \lambda w \in \mathcal{W}$  we conclude  $Tw - \lambda w = 0$  and  $Tu - \lambda u = 0$ . Since  $u \neq 0$  or  $w \neq 0$ , we have  $\lambda \in \sigma(T|_{\mathcal{U}})$  or  $\lambda \in \sigma(T|_{\mathcal{W}})$ .

Assume  $\lambda \in \sigma(T)$  and  $\lambda \notin \sigma(T|_{\mathcal{W}})$ . Then  $\mathcal{N}(T - \lambda I) \cap \mathcal{W} = \{0\}$ . Lemma 2.4 applies to the operator  $T - \lambda I$  and yields (10). Since  $\lambda \in \sigma(T)$ ,  $\mathcal{N}(T - \lambda I) \neq \{0\}$ . Now (10) with  $j = 1$  yields  $\lambda \in \sigma(T|_{\mathcal{U}})$ .  $\square$

dect

**Theorem 2.6.** *Let  $\mathcal{V}$  be a finite dimensional vector space over  $\mathbb{C}$ ,  $n = \dim \mathcal{V}$  and let  $T \in \mathcal{L}(\mathcal{V})$ . Let  $\lambda_1, \dots, \lambda_k$ , be all the distinct eigenvalues of  $T$ . Set*

$$\mathcal{W}_j = \mathcal{N}((T - \lambda_j I)^n) \quad \text{and} \quad \dim \mathcal{W}_j = n_j, \quad j \in \{1, \dots, k\}.$$

Then

dtb

(a) Each of the subspaces  $\mathcal{W}_1, \dots, \mathcal{W}_k$ , is invariant under  $T$ .

dtb

(b)  $\mathcal{V} = \mathcal{W}_1 \oplus \dots \oplus \mathcal{W}_k$ .

dtb

(c) Set  $T_j = T|_{\mathcal{W}_j}$  and  $N_j = T_j - \lambda_j I$ ,  $j \in \{1, \dots, k\}$ . Then  $N_j^{n_j} = 0$ , that is,  $N_j$  is a nilpotent operator on  $\mathcal{W}_j$ .

*Proof.* (a) Since  $T$  commutes with each of the operators  $(T - \lambda_j I)^d$ ,  $j \in \{1, \dots, k\}$  Lemma 2.1 implies that each subspace  $\mathcal{W}_1, \dots, \mathcal{W}_k$ , is an invariant subspace of  $T$ .

To prove (b) we proceed by mathematical induction on the number  $k$  of distinct eigenvalues of  $T$ . We first prove the base step. Assume that  $\lambda$  is the only eigenvalue of  $T$ . Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  be a basis of  $\mathcal{V}$  such that the matrix  $M_{\mathcal{B}}^{\mathcal{B}}(T)$  is upper triangular. Then, as we proved earlier all the diagonal entries of  $M_{\mathcal{B}}^{\mathcal{B}}(T)$  equal to  $\lambda$ . From the definition of  $M_{\mathcal{B}}^{\mathcal{B}}(T)$  it follows that

$$(T - \lambda I)(\text{span}\{v_1, \dots, v_j\}) \subseteq (\text{span}\{v_1, \dots, v_{j-1}\}) \quad \text{for all } j \in \{2, \dots, n\}.$$

Therefore

$$\begin{aligned} (T - \lambda I)^n(\mathcal{V}) &= (T - \lambda I)^{n-1}(T - \lambda I)(\text{span}\{v_1, \dots, v_n\}) \\ &\subseteq (T - \lambda I)^{n-1}(\text{span}\{v_1, \dots, v_{n-1}\}) \\ &\quad \vdots \\ &\subseteq (T - \lambda I)(T - \lambda I)(\text{span}\{v_1, v_2\}) \\ &\subseteq (T - \lambda I)(\text{span}\{v_1\}) \\ &= \{0_{\mathcal{V}}\}. \end{aligned}$$

Thus  $\mathcal{V} = \mathcal{N}((T - \lambda I)^n)$ . This completes the proof of the base case.

Now we prove the inductive step. Let  $k \in \mathbb{N}$  and assume that the statement is true for an operator with  $k$  distinct eigenvalues. Let  $T$  be an operator with  $k+1$  distinct eigenvalues  $\lambda_1, \dots, \lambda_k, \lambda_{k+1}$ . For convenience we set  $\lambda_{k+1} = \lambda$ . Then, by assumption  $\lambda \neq \lambda_j$  for all  $j \in \{1, \dots, k\}$ . We set

$$\mathcal{U} = \mathcal{R}((T - \lambda I)^n) \quad \text{and} \quad \mathcal{W} = \mathcal{N}((T - \lambda I)^n).$$

Since  $T$  and  $(T - \lambda I)^n$  commute, Lemma 2.1 implies that both  $\mathcal{U}$  and  $\mathcal{W}$  are invariant under  $T$ .

Next we prove that

$$\mathcal{R}((T - \lambda I)^n) \cap \mathcal{N}((T - \lambda I)^n) = \mathcal{U} \cap \mathcal{W} = \{0\}. \quad (11) \quad \boxed{\text{eq-triv}}$$

(Prove this as an exercise.)

By the Rank-Nullity theorem

$$\mathcal{V} = \mathcal{U} \oplus \mathcal{W}. \quad (12) \quad \boxed{\text{eq-ds}}$$

(Provide details as an exercise.)

By Corollary 2.3

$$\mathcal{N}((T - \lambda_j I)^n) \subseteq \mathcal{U} \quad \text{for all } j \in \{1, \dots, k\}. \quad (13) \quad \boxed{\text{eq-incl}}$$

Let  $m = \dim \mathcal{U}$ . Denote by  $S$  the restriction of  $T$  onto  $\mathcal{U}$ . The inclusion in (13) implies that  $\lambda_1, \dots, \lambda_k$  are eigenvalues of  $S$ . Similarly, (11) implies that  $\lambda$  is not an eigenvalue of  $S$ . Now Corollary 2.5 yields

$$\sigma(S) = \{\lambda_1, \dots, \lambda_k\}.$$

The second claim of Corollary 2.5 implies

$$\mathcal{N}((T - \lambda_j I)^n) = \mathcal{N}((S - \lambda_j I)^n).$$

Since  $n > m = \dim \mathcal{U}$  we have

$$\mathcal{N}((S - \lambda_j I)^m) = \mathcal{N}((S - \lambda_j I)^{m+1}) = \dots = \mathcal{N}((S - \lambda_j I)^n).$$

Therefore,

$$\mathcal{N}((T - \lambda_j I)^n) = \mathcal{N}((S - \lambda_j I)^m). \quad (14) \quad \boxed{\text{eq-ges=ges}}$$

The inductive hypothesis applies to  $S$ . Therefore

$$\mathcal{U} = \mathcal{R}((T - \lambda I)^n) = \bigoplus_{j=1}^k \mathcal{N}((S - \lambda_j I)^m). \quad (15) \quad \boxed{\text{eq-ihc}}$$

Now (15), (14), and (12) yield

$$\mathcal{V} = \bigoplus_{j=1}^{k+1} \mathcal{N}((T - \lambda_j I)^n).$$

Now we prove (c). Lemma 2.1 implies that  $\mathcal{W}_j$  is an invariant subspace of  $T - \lambda_j I$ . Denote by  $N_j$  the restriction of  $T - \lambda_j I$  to its invariant subspace  $\mathcal{W}_j$  and by  $T_j$  the restriction of  $T$  to  $\mathcal{W}_j$ . Then,  $T_j = \lambda_j I + N_j$  and the operator  $N_j$  is nilpotent.  $\square$

$\boxed{\text{dcp}}$

**Definition 2.7.** Let  $k \in \{1, \dots, n\}$  be such that  $\lambda_1, \dots, \lambda_k$  are all the distinct eigenvalues of  $T$ . Set

$$n_j = \dim \mathcal{N}((T - \lambda_j)^n), \quad j \in \{1, \dots, k\}.$$

The number  $n_j$  is called the *algebraic multiplicity* of the eigenvalue  $\lambda_j$ . The polynomial

$$p(z) = (z - \lambda_1)^{n_1} \dots (z - \lambda_k)^{n_k} \quad (16) \quad \boxed{\text{eqcnp2}}$$

is called the *characteristic polynomial* of  $T$ .

### 3 The Jordan Normal Form

Let  $T$  be an operator on a vector space  $\mathcal{V}$  over  $\mathbb{C}$ . Let  $\lambda$  be an eigenvalue of  $T$  and let  $v$  be such that  $(T - \lambda I)^l v = 0_{\mathcal{V}}$  and  $(T - \lambda I)^{l-1} v \neq 0_{\mathcal{V}}$ . Then the system of vectors

$$(T - \lambda I)^{l-1} v, (T - \lambda I)^{l-2} v, \dots, (T - \lambda I)v, v, \quad (17) \quad \boxed{\text{jca1}}$$

is called a *Jordan chain* of  $T$  corresponding to the eigenvalue  $\lambda$ . The vectors in (17) are called *generalized eigenvectors* (or *root vectors*) corresponding to the eigenvalue  $\lambda$ .

Let  $\mathcal{W}$  be a subspace of  $\mathcal{V}$  generated by a Jordan chain

$$v_j = (T - \lambda I)^{l-j} v, \quad j \in \{1, \dots, l\},$$

of  $T$ . Note that the vector  $v_1 = (T - \lambda I)^{l-1}v$  is an eigenvector of  $T$  corresponding to the eigenvalue  $\lambda$ . Therefore  $Tv_1 = \lambda v_1$ . We also have

$$Tv_j = (T - \lambda I)v_j + \lambda v_j = v_{j-1} + \lambda v_j, \quad j \in \{1, \dots, l\}.$$

It follows that  $\mathcal{W}$  is an invariant subspace of  $T$ . If we denote by  $A$  the restriction of  $T$  to  $\mathcal{W}$ , then the matrix representation of  $A$  with respect to the basis  $\{v_1, \dots, v_l\}$  is

$$\begin{bmatrix} \lambda & 1 & 0 & \cdots & 0 & 0 \\ 0 & \lambda & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & \cdots & \lambda & 1 \\ 0 & 0 & 0 & \cdots & 0 & \lambda \end{bmatrix}. \quad (18) \quad \boxed{\text{jbl1}}$$

A matrix of this form is called a *Jordan block* corresponding to the eigenvalue  $\lambda$ . In words: a Jordan block corresponding to the eigenvalue  $\lambda$  is a square matrix with all elements on the main diagonal equal to  $\lambda$  and all elements on the superdiagonal equal to 1.

A basis for  $\mathcal{V}$  which consists of Jordan chains of  $T$  is called a *Jordan basis* for  $\mathcal{V}$  with respect to  $T$ .

If a basis  $\mathcal{B}$  for  $\mathcal{V}$  is a Jordan basis with respect to  $T$  then the matrix  $M_{\mathcal{B}}(T)$  has Jordan blocks of different sizes on the diagonal and all other elements of  $M_{\mathcal{B}}(T)$  are zeros. Each eigenvalue of  $T$  is represented in  $M_{\mathcal{B}}(T)$  by one or more Jordan blocks;

$$\begin{bmatrix} \boxed{\begin{matrix} \lambda_1 & 1 & \cdots & 0 \\ 0 & \lambda_1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \\ 0 & 0 & \cdots & \lambda_1 \end{matrix}} & \begin{matrix} 0 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \end{matrix} \\ \begin{matrix} 0 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \end{matrix} & \boxed{\begin{matrix} \lambda_2 & 1 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \\ 0 & 0 & \cdots & \lambda_2 \end{matrix}} \\ \cdots & \cdots \end{bmatrix}. \quad (19) \quad \boxed{\text{jnf}}$$

In the above matrix  $\lambda_1$  and  $\lambda_2$  are not necessarily distinct eigenvalues. A matrix of the form (19) is called the *Jordan normal form* for  $T$ . More precisely, a square matrix  $M = [a_{j,k}]$  is a *Jordan normal form* for  $T$  if:

- (i) all elements of  $M$  outside of the main diagonal and the superdiagonal are 0,

- (ii) all elements on the main diagonal of  $M$  are eigenvalues of  $T$ ,
- (iii) all elements on the superdiagonal of  $M$  are either 1 or 0, and,
- (iv) if  $a_{j-1,j-1} \neq a_{j,j}$ , with  $j \in \{2, \dots, n\}$ , then  $a_{j-1,j} = 0$ .

**Theorem 3.1.** *Let  $\mathcal{V}$  be a vector space over  $\mathbb{C}$  and let  $T$  be a linear operator on  $\mathcal{V}$ . Then  $\mathcal{V}$  has a Jordan basis with respect to  $T$ .*

*Proof.* We use the notation and the results of Theorem 2.6. Let  $j \in \{1, \dots, k\}$ . It is important to notice that each Jordan chain of the nilpotent operator  $N_j$  is a Jordan chain of  $T$  which corresponds to the eigenvalue  $\lambda_j$ . Since  $N_j$  is a nilpotent operator in  $\mathcal{L}(\mathcal{W}_j)$ , by Theorem 1.1 there exists a basis  $\mathcal{B}_j = \{v_{j,1}, \dots, v_{j,n_j}\}$  for  $\mathcal{W}_j$  which consists of Jordan chains of  $N_j$ . Consequently,  $\mathcal{B}_j$  consists of Jordan chains of  $T$ . Since  $\mathcal{V}$  is a direct sum of  $\mathcal{W}_1, \dots, \mathcal{W}_k$ , the union  $\mathcal{B} = \mathcal{B}_1 \cup \dots \cup \mathcal{B}_k$ , that is,

$$\mathcal{B} = \{v_{1,1}, \dots, v_{1,n_1}, v_{2,1}, \dots, v_{2,n_2}, \dots, v_{k,1}, \dots, v_{k,n_k}\}$$

is a basis for  $\mathcal{V}$ . This basis consists of Jordan chains of  $T$ .

The matrix  $M_{\mathcal{B}}(T)$  is a block diagonal with the blocks  $M_{\mathcal{B}_j}(T_j)$ ,  $j \in \{1, \dots, k\}$ , on the diagonal and with zeros every where else:

$$M_{\mathcal{B}}^{\mathcal{B}}(T) = \begin{bmatrix} M_{\mathcal{B}_1}^{\mathcal{B}_1}(T_1) & 0 & \dots & 0 \\ 0 & M_{\mathcal{B}_2}^{\mathcal{B}_2}(T_2) & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & M_{\mathcal{B}_k}^{\mathcal{B}_k}(T_k) \end{bmatrix}.$$

Since  $T_j = \lambda_j I + N_j$ , we have

$$M_{\mathcal{B}_j}(T_j) = \lambda_j I + M_{\mathcal{B}_j}(N_j).$$

Thus all the elements on the main diagonal of  $M_{\mathcal{B}_j}^{\mathcal{B}_j}(T_j)$  equal  $\lambda_j$  and all the elements of superdiagonal of  $M_{\mathcal{B}_j}^{\mathcal{B}_j}(T_j)$  are either 1 or 0. If there are exactly  $m_j$  Jordan chains in the basis  $\mathcal{B}_j$ , then 0 appears exactly  $m_j - 1$  times on the superdiagonal of  $M_{\mathcal{B}_j}^{\mathcal{B}_j}(T_j)$ . Therefore  $M_{\mathcal{B}}^{\mathcal{B}}(T)$  is a Jordan normal form for  $T$ .  $\square$